



UTSW/BioTel EMS TRAINING BULLETIN

October 2014

EMS TB 14-006

Ebola Virus Disease (EVD)

Purpose:

1. To inform & provide management recommendations to UTSW/BioTel EMS System EMS Providers about Ebola Virus Disease (**EVD**).

Background:

1. A patient with **EVD**, also known as Ebola Hemorrhagic Fever, was recently evaluated & transported to Texas Health Presbyterian Hospital by Dallas Fire-Rescue paramedics.

EVD Transmission:

1. Ebola virus can be transmitted **ONLY** via **DIRECT** contact with the bodily fluids (e.g. blood, saliva, emesis, feces, urine or semen) of a **SYMPTOMATIC, INFECTED** person.
2. Ebola virus **CANNOT** be transmitted by casual contact, such as being in the same room with a patient.
3. Persons infected with Ebola virus are **ONLY** contagious when they are **SYMPTOMATIC**. In other words, patients infected with Ebola virus are **NOT** contagious during the asymptomatic incubation period immediately following infection (which ranges from 2 to 21 days, with an average 8 to 10 days).

Patient Signs and Symptoms of EVD that suggest the possibility that a patient may be potentially contagious:

1. Fever
2. Headache
3. Malaise
4. Body aches
5. Fatigue
6. Nausea
7. Vomiting
8. Diarrhea
9. Bleeding or bruising of unknown cause

Patient Evaluation Procedures and Precautions:

1. EMS Providers who evaluate ANY patient with fever **AND** any of the flu-like symptoms listed above shall ALWAYS use STANDARD, DROPLET & CONTACT Precautions.
2. OTHER HISTORY factors associated with risk of exposure for which STANDARD, DROPLET & CONTACT precautions are indicated during patient evaluation/transport:
 - a. The **ASYMPTOMATIC INCUBATION PERIOD** after exposure to Ebola virus ranges from 2 to 21 days. Therefore, even though the patient may not be symptomatic (or contagious) during this period, any patient who has had contact with a person with EVD should be considered to be at risk.
 - b. **TRAVEL HISTORY:** Any patient who has recently traveled to the DFW Metroplex from West Africa should be considered to be at risk.
3. EMS Providers shall follow agency-specific guidelines for notification of EMS Chain of Command after evaluation and/or transport of a patient with the symptoms listed above.
4. In addition, BioTel and/or the receiving hospital **shall be notified** before or during transport of such patients, so that the receiving hospital Emergency Department staff can be notified prior to arrival.
5. Finally, any and all concerns regarding the possibility of EVD shall be communicated directly to the receiving hospital Emergency Department staff immediately upon arrival.

Disinfection and Decontamination Procedures/Precautions:

1. Ebola virus is very fragile. It cannot easily survive outside of body fluids. Therefore, disinfection is relatively simple.
2. Soap and water and standard disinfecting/cleaning solutions readily kill the virus.
3. Standard, Droplet and Contact precautions (including the use of standard personal protective equipment (PPE)), disposal of contaminated items, and equipment cleaning are sufficient to prevent exposure. EMS Providers should follow agency-specific procedures for routine decontamination of the vehicle and equipment.
4. Procedures with high risk of aerosolization of body fluids (e.g. airway suctioning and advanced airway management) require use of an N95 respirator or higher.

Local, County, State and Federal Collaboration:

1. UTSW/BioTel Fire/EMS agency command staff members are working collaboratively with all local, County, State and Federal colleagues, including local hospital partners, Parkland BioTel, City governments, Dallas County Health & Human Services, the Centers for Disease Control and Prevention (CDC), & the UTSW/BioTel EMS Medical Direction Team to provide timely, accurate information to all EMS Providers during this challenging period.

FINAL NOTES:

- **Additional updates will be provided as new information and recommendations become available.**
- **EMS Providers are encouraged to contact their EMS Supervisor or BioTel at any time with questions or concerns.**

Ebola Hemorrhagic Fever

Ebola hemorrhagic fever (Ebola HF) is one of numerous Viral Hemorrhagic Fevers. It is a severe, often fatal disease in humans and nonhuman primates (such as monkeys, gorillas, and chimpanzees).

Ebola HF is caused by infection with a virus of the family *Filoviridae*, genus *Ebolavirus*. When infection occurs, symptoms usually begin abruptly. The first Ebolavirus species was discovered in 1976 in what is now the Democratic Republic of the Congo near the Ebola River. Since then, outbreaks have appeared sporadically.

There are five identified subspecies of *Ebolavirus*. Four of the five have caused disease in humans: Ebola virus (*Zaire ebolavirus*); Sudan virus (*Sudan ebolavirus*); Taï Forest virus (*Taï Forest ebolavirus*, formerly *Côte d'Ivoire ebolavirus*); and Bundibugyo virus (*Bundibugyo ebolavirus*). The fifth, Reston virus (*Reston ebolavirus*), has caused disease in nonhuman primates, but not in humans.

The natural reservoir host of ebolaviruses remains unknown. However, on the basis of available evidence and the nature of similar viruses, researchers believe that the virus is zoonotic (animal-borne) with bats being the most likely reservoir. Four of the five subtypes occur in an animal host native to Africa.

A host of similar species is probably associated with Reston virus, which was isolated from infected cynomolgous monkeys imported to the United States and Italy from the Philippines. Several workers in the Philippines and in US holding facility outbreaks became infected with the virus, but did not become ill.

Transmission

Because the natural reservoir of ebolaviruses has not yet been proven, the manner in which the virus first appears in a human at the start of an outbreak is unknown. However, researchers have hypothesized that the first patient becomes infected through contact with an infected animal.

When an infection does occur in humans, there are several ways in which the virus can be transmitted to others. These include:

- direct contact with the blood or secretions of an infected person
- exposure to objects (such as needles) that have been contaminated with infected secretions

The viruses that cause Ebola HF are often spread through families and friends because they come in close contact with infectious secretions when caring for ill persons.

During outbreaks of Ebola HF, the disease can spread quickly within health care settings (such as a clinic or hospital). Exposure to ebolaviruses can occur in health care settings where hospital staff are not wearing appropriate protective equipment, such as masks, gowns, and gloves.

Proper cleaning and disposal of instruments, such as needles and syringes, is also important. If instruments are not disposable, they must be sterilized before being used again. Without adequate sterilization of the instruments, virus transmission can continue and amplify an outbreak.

Signs and Symptoms

Symptoms of Ebola HF typically include:

- Fever
- Headache
- Joint and muscle aches
- Weakness
- Diarrhea
- Vomiting
- Stomach pain
- Lack of appetite

Some patients may experience:

- A Rash
- Red Eyes
- Hiccups
- Cough
- Sore throat
- Chest pain
- Difficulty breathing
- Difficulty swallowing
- Bleeding inside and outside of the body

Symptoms may appear anywhere from 2 to 21 days after exposure to ebolavirus though 8-10 days is most common.

Some who become sick with Ebola HF are able to recover, while others do not. The reasons behind this are not yet fully understood. However, it is known that patients who die usually have not developed a significant immune response to the virus at the time of death.

Risk of Exposure

In Africa, confirmed cases of Ebola HF have been reported in:

- Guinea
- Liberia
- Sierra Leone
- Democratic Republic of the Congo (DRC)
- Gabon
- South Sudan
- Ivory Coast
- Uganda
- Republic of the Congo (ROC)
- South Africa (imported)

The natural reservoir host of ebolaviruses, and the manner in which transmission of the virus to humans occurs, remain unknown. This makes risk assessment in endemic areas difficult. Currently, all cases of illness or death have occurred in Africa; no case has been reported in the United States.

During outbreaks of Ebola HF, those at highest risk include health care workers and the family and friends of an infected individual. Health care workers in Africa should consult the Infection Control for Viral Hemorrhagic Fevers In the African Health Care Setting to learn how to prevent and control infections in these setting. Medical professionals in the United States should consult the Interim Guidance for Managing Patients with Suspected Viral Hemorrhagic Fever in U.S. Hospitals.

Diagnosis

Diagnosing Ebola HF in an individual who has been infected for only a few days is difficult, because the early symptoms, such as red eyes and a skin rash, are nonspecific to ebolavirus infection and are seen often in patients with more commonly occurring diseases.

However, if a person has the early symptoms of Ebola HF and there is reason to believe that Ebola HF should be considered, the patient should be isolated and public health professionals notified. Samples from the patient can then be collected and tested to confirm infection.

Laboratory tests used in diagnosis include:

Timeline of Infection	Diagnostic tests available
Within a few days after symptoms begin	- Antigen-capture enzyme-linked immunosorbent assay (ELISA) testing - IgM ELISA - Polymerase chain reaction (PCR) - Virus isolation
Later in disease course or after recovery	- IgM and IgG antibodies
Retrospectively in deceased patients	- Immunohistochemistry testing - PCR - Virus isolation

Treatment

Standard treatment for Ebola HF is still limited to supportive therapy. This consists of:

- balancing the patient's fluids and electrolytes
- maintaining their oxygen status and blood pressure
- treating them for any complicating infections

Timely treatment of Ebola HF is important but challenging since the disease is difficult to diagnose clinically in the early stages of infection. Because early symptoms such as headache and fever are nonspecific to ebolaviruses, cases of Ebola HF may be initially misdiagnosed.

However, if a person has the early symptoms of Ebola HF and there is reason to believe that Ebola HF should be considered, the patient should be isolated and public health professionals notified. Supportive therapy can continue with proper protective clothing until samples from the patient are tested to confirm infection.

Experimental treatment has been tested and proven effective in animal models but has not yet been used in humans.

Prevention

The prevention of Ebola HF presents many challenges. Because it is still unknown how exactly people are infected with Ebola HF, there are few established primary prevention measures.

When cases of the disease do appear, there is increased risk of transmission within health care settings. Therefore, health care workers must be able to recognize a case of Ebola HF and be ready to employ practical viral hemorrhagic fever isolation precautions or barrier nursing techniques. They should also have the capability to request diagnostic tests or prepare samples for shipping and testing elsewhere.

Barrier nursing techniques include:

- wearing of protective clothing (such as masks, gloves, gowns, and goggles)
- the use of infection-control measures (such as complete equipment sterilization and routine use of disinfectant)
- isolation of Ebola HF patients from contact with unprotected persons

The aim of all of these techniques is to avoid contact with the blood or secretions of an infected patient. If a patient with Ebola HF dies, it is equally important that direct contact with the body of the deceased patient be prevented.

CDC, in conjunction with the World Health Organization, has developed a set of guidelines to help prevent and control the spread of Ebola HF. Entitled Infection Control for Viral Hemorrhagic Fevers In the African Health Care Setting, the manual describes how to:

- recognize cases of viral hemorrhagic fever (such as Ebola HF)
- prevent further transmission in health care setting by using locally available materials and minimal financial resources